

STUDY ON ARSENIC LEVEL IN GROUND WATER OF DELHI USING HYDRIDE GENERATOR ACCESSORY COUPLED WITH ATOMIC ABSORPTION SPECTROPHOTOMETER

Sanjeev Lalwani*, T.D. Dogra*, D.N. Bhardwaj*, R.K. Sharma*, O.P. Murty* and Aarti Vij**

**Department of Forensic Medicine, **Dept. of Hospital Administration, All India Institute of Medical Sciences, Ansari Nagar, New Delhi 110 029*

ABSTRACT

Surveillance of drinking water is essentially a health measure intended to protect the public from water borne diseases. Hydride generator accessory coupled with atomic absorption spectrophotometer was used to analyze arsenic level in 49 ground water samples collected from different areas of Delhi. Arsenic level in ground water samples was in the range of 0.0170 to 0.100 ppm (Mean-0.0431, Standard Deviation-0.0136, Std. error of Mean-0.00194) with minimum concentration at Raney Well No. 7 (0.0170 ppm) and maximum at Kotla Mubarak Pur (0.100 ppm). Arsenic containing sediments and percolation of chemicals into soil as the result of dumping of garbage rich in chemicals into open landfills could be the possible source of arsenic in ground water of Delhi. Extensive survey and continuous monitoring is required to be made to assess the magnitude of problem and earlier intervention.

KEY WORDS

Arsenic, ground water, hydride generator accessory, atomic absorption spectrophotometer.

INTRODUCTION

Arsenic is a ubiquitous, nonessential, bio-accumulative element known since 387 B.C. Its compounds have been used as medicine, insecticides, wood preservatives, dye and industrial chemicals (1). Toxic exposure of human beings to arsenic and its compounds may occur through air, water, seafood, tobacco smoking, beverages medicinal preparations and industrial chemicals (2). Worldwide, the main reason for a chronic human intoxication with arsenic is the intake of contaminated drinking water (3).

Arsenic and its compounds are naturally present in low concentration at places with high geothermal activities (2). The current drinking water quality guideline by WHO for arsenic is 0.01 ppm (4). Long term exposure to elevated level of arsenic in

water may cause serious health hazards like rhinopharyngitis, pulmonary insufficiency, interstitial fibrosis, hyperkeratosis of palm and soles, melanosis, noncirrhotic portal fibrosis, myocardial damage with ECG changes, hypertension, peripheral vascular disease, sensorimotor polyneuropathy, retrobulbar neuritis, encephalopathy, bilateral optic atrophy, deafness, bowens disease, haemangioen-dothelioma of liver, leukemia, malignancy of stomach, urinary bladder, oesophagous, kidney, bone and lungs.(2, 5-15)

In India, natural exposure of man to arsenic through drinking water of wells, hand pumps and springs in Chandigarh and its surrounding areas was first highlighted in 1976 (16-17). Arsenic contamination of ground water in West Bengal was first reported in 1983, but the relationship of dermatological manifestation to arsenic in water of tube wells of West Bengal was established by 1984 (18). However, the magnitude of the problem remains undetermined till 1995. So far more than 3,00,000 people of 9 districts of West Bengal have developed arsenicosis due to consumption of arsenic contaminated drinking water (19). Toxic level of arsenic in ground water has also been reported from villages of Rajnandgaon district of Madhya Pradesh (19) and some areas of Bihar (20). 1n

Author for correspondence :

Dr.Sanjeev Lalwani

C/o Dr. D.N.Bhardwaj

E-14, AIIMS Ansari Nagar

New Delhi-110 029, INDIA

E mail : drsalal@rediffmail.com

sanjulalwani2001@yahoo.com

1997, people of 34 districts of Bangladesh were reported to be drinking arsenic contaminated water (6). In the combined area of West Bengal and Bangladesh about 150 million people are at risk from arsenic contaminated ground water (21). Arsenic contamination of well water in Minnetosa and Taiwan has caused signs and symptoms of arsenic toxicity (22, 23). Similar results have been reported from Northern Mexico, Nova Scotia Canada, Finland, Alaska and China (24-28).

In Delhi, drinking water supply is not from single source. Sixty eight percent of the population is getting supply from water of Yamuna River. Ground water sources like tube wells, hand pumps and borings are the other sources. As far as quality is concerned there is regular monitoring of physical parameters only. Suspended impurities such as metals are not monitored. Most of the times ground water is directly used. The present study was aimed to analyze arsenic level in ground water sources of Delhi and to look for the possible source of contamination.

MATERIALS AND METHODS

Sampling

Water samples were collected by using plastic bottles of capacity of one litre. All bottles used were washed with 1% nitric acid (BDH, E.Merck(India) Ltd, Mumbai (Maharashtra), India). Before collecting the sample, bottles were rinsed three-five times with the water to be filled. After filling with water few drops of concentrated nitric acid were added to the water samples for preservation till analysis. Total 49 ground water samples involving tube wells, hand pumps and borings were collected from different areas of Delhi as shown in Table 1.

Preparation of Samples

Samples were filtered using 0.45 μ m Whatman filter paper. Initial filter was used for the rinsing of volumetric flask.

Standard

All the solutions were prepared with distil (deionized) water. Arsenic stock solution containing 980 μ gm per ml (980 ppm) (Sigma Chemical Company, St. Louis, USA) was used for preparing the standards.

METHODS

Arsenic in water was determined according to the study of C.J. Wyatt *et al.* (15). Air -Acetylene flame type of Atomic Absorption Spectrophotometer (AAS 4127) along with Hydride Generator Accessory both

of ECIL Company were used for analysis of arsenic in water. Hollow Cathode Lamp of Arsenic was used (10 mA, wavelength 193.7). Nitrogen was used as inert gas. Sodium Borohydride (CDH Lab. Reagents, Mumbai-New Delhi, India) and Hydrochloric acid -HCl Min.35% (E. Merck (India) Ltd., Mumbai, Maharashtra, (India) were used for the hydride formation. Arsenic V was reduced to As III with a reduction with Potassium Iodide (Qualigens Fine Chemicals, Mumbai (Maharashtra), India) prior to the formation of the hydride as the sensitivity of instrument for detection of As III was higher 70-80% in comparison to As V (10-20%). Results were analysed using SPSS and Barlett's Chi Square test.

RESULTS

Arsenic level in 49 ground water samples collected from different areas of Delhi is shown in table 1. Arsenic level detected was in the range of 0.0170 ppm to 0.100 ppm (Mean-0.0431, Standard Deviation-0.0136, Std. error of Mean-0.00194). The minimum value was found in the ground water sample of Raney well no.7 (0.017 ppm) and maximum value in the ground water sample of Kotla Mubarak Pur (0.100 ppm). Arsenic level in tube wells of Bawana was found to be in the range of 0.022-0.044 ppm (Mean-0.034, Standard Deviation-0.011, Std. error of Mean-0.0065). In all the areas arsenic level in ground water is more than permissible limit of WHO (1993)(4). The areas where the arsenic level in ground water was more than 0.050 ppm were Masjid Moth, Gulmohar Park, Raney Well (p), Lajpat Nagar, Saket and Kotla Mubarak Pur.

DISCUSSION

The hydride generation technique with subsequent atomic absorption spectrophotometer in suitable flame as used in present study is a well known method for determination of arsenic. Simplicity and sensitivity of both the instrument in combination, best meet the requirement for economic and fast analysis of arsenic and other hydride forming element even if they are present in trace amount in environmental media and biological fluid (29). This technique have been used in various studies reported in literature (5-7, 15, 24).

Arsenic level in 49 ground water samples collected from different areas of Delhi was in the range of 0.0170 to 0.1000 ppm, which is more than WHO recommended value for arsenic in drinking water (0.01ppm). Recent study conducted in Department of Environmental science in Guru Govind Singh Indraprastha University of Delhi has also revealed that concentration of contaminants like arsenic in

Table 1. Showing arsenic level in ground water samples of Delhi

S.No.	Source	No. of Samples	Arsenic Level (PPm)			
			Range	Mean	S.D.	SEM
1	Bawana	3	0.022-0.044	0.034	0.011	0.006
2	Masjid Moth	1	0.052	0.052	0	0
3	Gulmohar Park	1	0.064	0.064	0	0
4	Malviya Nagar	2	0.034-0.049	0.043	0.013	0.009
5	Saket	1	0.052	0.052	0	0
6	Mehrauli	3	0.040-0.049	0.044	0.005	0.003
7	Lado Sarai	1	0.042	0.042	0	0
8	S.J.Enclave	2	0.022	0.022	0	0
9	Green Park	1	0.031	0.031	0	0
10	Lajpat Nagar	1	0.057	0.057	0	0
11	K M Pur	1	0.100	0.100	0	0
12	Janakpuri	1	0.032	0.032	0	0
13	Moti Bagh	4	0.036-0.056	0.043	0.004	0.002
14	Satya Niketan	1	0.032	0.032	0	0
15	Sagarpur	1	0.036	0.036	0	0
16	RK Puram-7	2	0.050	0.050	0	0
17	RK Puram-3	1	0.050	0.050	0	0
18	RK Puram-6	3	0.042-0.050	0.047	0.005	0.003
19	Raney Well P	1	0.064	0.064	0	0
20	Raney well V	1	0.048	0.048	0	0
21	Tilak Nagar	7	0.032-0.048	0.036	0.006	0.002
22	Khyala	2	0.036-0.060	0.048	0.016	0.012
23	Raghubir Nagar	3	0.036-0.050	0.043	0.007	0.004
24	Raney Well Alipur	1	0.034	0.034	0	0
25	Raney Well 7	1	0.017	0.017	0	0
26	Okhla	1	0.044	0.044	0	0
27	East Patel Nagar	1	0.046	0.046	0	0
28	West Patel Nagar	1	0.053	0.053	0	0
Total no. of Samples		= 49				
Mean Value of Arsenic		= 0.0431				
Std. Deviation		= 0.0136				
Std. Error of Mean		= 0.002				

the ground water of Delhi exceeds the permissible limits (30).

The mechanism of presence of arsenic in ground water is not clearly known. Nickson et al suggested anoxic reduction of arseniferous iron-oxyhydroxides as the possible source of arsenic in ground water (31-32). However, this theory does not explain progressively increasing arsenic level in tube wells (33-34). According to other theory suggested by Chakraborti *et al.* on the basis of sediment analysis high volume extraction of ground water causes exposure of deltaic segments to air which through oxidation reaction causes decomposition of iron pyrites. Arsenic is released in the process which then oxidised into arsenite and arsenate both of which are soluble in ground water (32-33). Le Blancs suggested catalytic oxidation of metal sulfide by stromatolytique micro organism as the possible mechanism (32).

Chakraborti *et al.* has reported elevated levels of iron in ground water samples of Bihar (20). Similarly Nag *et al.* (35) reported high concentration of iron in ground water samples in Purbasthali, Burdwan (West Bengal). These studies favour the theory of Chakraborti et al as the possible mechanism of presence of arsenic and iron in ground water of West Bengal and Bihar. Gallagher et al has mentioned that stability of As (III) and Arsenic (V) in iron rich drinking water can be affected by formation of iron precipitates (Iron Hydroxides and Oxides) (36). In Delhi, to meet the increasing public demand of water there is uncontrolled and excessive extraction of ground water. In an unofficial statement one of the Raney well used as water source was said to contain high iron level. Goswami *et al.* has reported elevated iron level in ground water of Delhi (37). Thus the presence of arsenic in ground water of Delhi can also be explained by the theory of Chakraborti et al. Further, everyday dumping of thousands of tons of garbage containing hazardous chemicals in open landfills of Delhi may be adding arsenic to the ground water (38). Chakraborti *et al.* (7) has reported the contamination of ground water due to discharge of effluent from the pesticide manufacturing industry in Calcutta. High porosity of soil causes percolation of chemicals and contamination of ground water.

In all the areas arsenic level in ground water is more than permissible limit of WHO (1993) (4). The condition may deteriorate further particularly in view of present days continuous, uncontrolled and heavy extraction of ground water and excessive use of chemicals along with discharge of hazardous waste in landfills. Therefore, extensive survey and continuous monitoring is required to be made to

assess the magnitude of problem and earlier intervention.

REFERENCES

1. Subramanyam, B.V. (2000) Inorganic Irritant Poisons- Arsenic In: Modi's Textbook of Medical Jurisprudence and Toxicology, 22nd edn. Butterworths India Ltd, New Delhi, 103-127.
2. WHO (1981) International Programme on Chemical Safety (IPCS)- Environmental Health Criteria-18, Arsenic, World Health Organization, Geneva.
3. Gebel, T. (2000) Confounding variables in the environmental toxicology of arsenic. *Toxicology*, 144,155-162.
4. WHO (1993). *Guidelines for drinking water quality : Recommendations*, Vol. 1 2nd edn., World Health Organization Geneva.
5. Mandal, B.K., Chowdhury, T.R., Samanta, G., Basu, G.K., Chowdhury, P.P., Chanda, C.R., Lodh, D., Karan, N.K., Dhar, R.K., Tamili, D.K., Das, D., Saha, K.C. and Chakraborti D. (1996) Arsenic in groundwater in seven districts of West Bengal, India - The Biggest arsenic calamity in the world. *Curr. Sci.* 72, 976-985.
6. Dhar, R.K., Biswas, B.K., Samanta, G., Mandal, B.K., Chakraborti, D., Roy, S., Jafar, A., Islam, A., Ara, G., Kabir, S., Khan, A.W., Ahmed, K.A. and Hadi, S.A., (1997) Ground Water arsenic calamity in Bangladesh. *Curr. Sci.* 73, 48-59.
7. Chakraborti, D., Samanta, G., Mandal, B.K., Chowdhury, T.R., Chanda, C.R., Biswas, B.K., Dhar, R.K., Basu, K. and Saha, K.C., (1998) Calcutta's Industrial pollution: Ground water arsenic contamination in a residual area sufferings of people due to industrial effluent discharge- An eight year study report. *Curr. Sci.* 74, 346-355.
8. Chen, C.J., Hseush, Y.M., Lai, M.S., Shyu, M.P., Chen, S.Y., Wu, M.M., Kuo, T.L. and Tai, T.Y. (1995) Increased prevalence of hypertension and long term arsenic exposure. *Hypertension*, 25, 53-60.
9. Enterline, P.E., Day, R. and Marsh, G.M., (1995) Cancers related to exposure to arsenic at a copper smelter. *Occup. Env. Med.*, 52, 28-32.
10. Chen, C.J., Chuang, Y.C., Lin, T.M., Wu, H.Y. and Yoh, S. (1985) Malignant neoplasm among residents of blackfoot disease endemic area in Taiwan- High artesian well water and cancer.

- Cancer Res. 45, 5895-5899.
11. Chen, C.J., Chuang, Y.C., Lin, T.M., Wu, H.Y. and Yoh, S. (1986) A retrospective study of malignant neoplasm of bladder, lung and liver in blackfoot disease endemic area in Taiwan. *Br. J. Cancer*, 53, 399-405.
 12. Jaerup, L., Pershagen, G. and Wall, S. (1989) Cumulative arsenic exposure and Lung cancer in smelter workers- A dose response study. *Am. J. Ind. Med.*, 15, 31-41.
 13. Battista, G., Bartoli, D., Iaia, T.E., Dini, F., Fiumalbi, C., Giglioli, S. and Valiani, M. (1996) Art glass workers and Sino nasal cancer: - report of three cases. *Am. J. Ind. Med.*, 30,31-35.
 14. Rahmann, M., Wingren, G. and Axelson, O. (1996) Diabetes Mellitus among Swedish art glass workers- an effect of arsenic exposure. *Scand. J. Work Env. Health*, 22, 146-149.
 15. Wyatt, C.J., Lopez, O.V., Olivas Acosta, R.T. and Oliva Mendez, R. (1998) Excretion of arsenic in urine of children 7-11 years, exposed to elevated levels of arsenic in city water supply in Hermosillo, Sonora, Mexico. *Env. Res.* 78, 19-24.
 16. Datta D.V. (1976) Arsenic and non cirrhotic portal hypertension (Letter). *Lancet* 1, 433.
 17. Datta, D.V. and Kaul, M.K. (1976) Arsenic content of drinking water in villages of Northern India. A concept of arsenicosis. *J. Assoc. Phys. Ind.* 24, 599-604.
 18. Saha, K.C. (1984) Melanokeratosis from arsenic contaminated tubewell water. *Ind. J. Dermat.* 29, 37-46.
 19. Chakraborti, D., Biswas, B.K., Chowdhury, T.R., Basu, G.K., Mandal, B.K., Chowdhury, U.K., Mukherjee S.C., Gupta, J.P., Chowdhury, S.R. and Rathore K.C. (1999) Arsenic groundwater contamination and sufferings of people in Rajnandgaon district, Madhya Pradesh, India. *Curr. Sci.* 77, 502-504.
 20. Chakraborti, D., Mukherjee, S.C., Pati, S., Sengupta, S.K., Rahman, M.M., Chowdhury, U.K., Lodh, D., Chanda, C.R., Chakraborti, A.K. and Basu, G.K. (2003) Arsenic groundwater contamination in Middle Ganga Plain, Bihar, India A Future Danger. *Env. Health Persp.* 111, 1194-1201.
 21. Rahmann, M.M., Chowdhury, U.K., Mukherjee, S.C., Mandal, B.K., Paul, K., Lodh, D. *et al.* (2001). Chronic arsenic toxicity in Bangladesh and West Bengal India. A review commentary. *J. Toxicol. Clin. Toxicol.* 39, 683-700.
 22. Tsang, W.P., Chu, H.M., How, S.W., Fong, J.M., Lin, C.S. and Yeh, S. (1968) Prevalence of skin cancer in an endemic areas of chronic arsenicism in Taiwan. *J. Natl. Cancer Inst.* 40, 453-462.
 23. Luo, Z.D., Zhang, Y.M., Ma, L., Zhang, G.Y., He, X., Wilson, R., Byrd, D.M., Griffith, J.G., Lai, S., He, L. *et al.* (1997) Chronic arsenicism and cancer in Inner Mongolia. Consequences of well water arsenic levels greater than 50microgram per ltr. In: *Arsenic exposure and Health effects*, Eds. Chappel WR, Abernathy CO, and Calderson RL, Chappmann and Hall, New York, p55-68.)
 24. Rosas, I., Belmont, R., Ammeinta, A. and Baez, A. (1999) Arsenic concentration in water soil forage and milk in Camara Languera, Mexico. *Water, Soil and Pollution.* 112, 133-149.
 25. Grantham, D.A. and Jones, J.F. (1977) Arsenic concentration in water of wells in Nova Scotia Canada. *J. Am. Water Works Assoc.* 69, 652-657.
 26. Harrington, J.M., Middaugh, J.P., Morse, D.L. and Houseworth, J. (1978) A Survey of a population exposed to high concentration of arsenic in well water in Fairbanks Alaska. *Am. J. Epidemiology.* 108, 377-385.
 27. Kurttio, P., Komulainen, H., Hakala, E., Kahalin, H. and Pekkanen, J.(1998) Urinary excretion of arsenic species after exposure to arsenic present in drinking water. *Arch. Env. Cont. Toxicol.* 34, 297-305.
 28. Zhang, L. and Chen, C. (1997) Geographic and exposure population of drinking water with high concentration of arsenic in China. *Wei. Sheng. Van. Jiu.* 26, 310-313.
 29. Thompson, K.C. and Thomerson, D.R. (1974) Atomic absorption studies on the determination of antimony, arsenic, bismuth, germanium, lead, selenium, tellurium and tin by utilizing the generation of covalent hydrides. *Analyst* 99, 595-601.
 30. Agarwal, R. and Wankade, K. (2003). <http://www.indiatogether.org/2003/jun/env-mercury.htm>
 31. Nickson, R., Mc Arthur, J.M., Ravenscrott, P., Burgess, W.G. and Ahmed K.S.. (1998) Mechanism of arsenic release of ground water, Bangladesh and West Bengal. *Appl. Geochem.* 16, 403-413.

32. Mai, T.T., Ph, O.M., Phan, Long. and Mekong, P.E. Forum. Ground water arsenic contamination : Can it happen in the Mekong Delta? A Vietnamese Perspectives. <http://www.mekongforum.org>
33. Chakraborti, D., Basu, G.K., Biswas, B.K., Chowdhury, U.K., Rahman, M.M., Paul, K. *et al.* (2001) Characterisation of arsenic bearing sediments in gangetic delta of West Bengal-India. In :Arsenic exposure and Health effects, Eds., Chappel, W.R., Abernathy, C.O. and Calderson, R.L., Elsevier Science, New York, 27-52.
34. Das, D., Samanta, G., Mandal, B.K., Chowdhury, R.T., Chanda, C.R., Chowdhury, P.P. *et al.* (1996) Arsenic in ground water of six districts of West Bengal, India. *Env. Geochem. Health.* 18, 5-15.
35. Nag, J.K., Balram, V., Rubio, R., Alberti, J. and Das, A.K. (1996) Inorganic arsenic species in ground water. A Case study from Purbasthali (Burdwan), India. *J. Trace Element Med. Biol.*, 10, 20-24.
36. Gallagher, P.A., Schwell C.A., Wei, X. and Creed, J.T. (2001) Speciation and preservation of inorganic arsenic in drinking water. Source using EDTA.Separation and ICP-MS detection. *Env. Mon.* 3, 371-376.
37. Goswami, A. and Singh, A.K. (2002) Enrichment of iron (III), cobalt (II), nickel (II), and copper (II) by solid-phase extraction with 1, 8-dihydroxyanthraquinone anchored to silica gel before their determination by flame atomic absorption spectrometry. *Anal. Bioanal. Chem.* 374, 554-560.
38. Earth Crash, Documenting the Collapse of a Dying Planet. New Delhi Facing Environmental Disaster As Uncontrolled Dumping of 1,000s of Tons of Garbage Every Day Spreads Disease, Contaminates Groundwater with Lead, Arsenic, Pesticides. <http://www.eces.org/articles/static/98264880068578.shtml>